arXiv:math/9807052v1 [math.QA] 10 Jul 1998

Line bundles on quantum spheres

Tomasz Brzeziński*[1] and Shahn Majid†[2]

*Department of Mathematics, University of York, Heslington, York YO10 5DD, England
† DAMTP, University of Cambridge, Cambridge CB3 9EW, England

Abstract. The (left coalgebra) line bundle associated to the quantum Hopf fibration of any quantum two-sphere is shown to be a finitely generated projective module. The corresponding projector is constructed and its monopole charge is computed. It is shown that the Dirac q-monopole connection on any quantum two-sphere induces the Grassmannian connection built with this projector.

INTRODUCTION

In the standard approach to non-commutative geometry [7] (see [11] for an accessible introduction) the notion of a vector bundle is identified with that of a finitely generated projective module E of an algebra B. Recall that E is a finitely generated projective left B-module if E is isomorphic to $B^n e := \{(b_1, \dots, b_n)e \mid b_1, \dots, b_n \in B\}$, where e is an $n \times n$ matrix with entries from B such that $e^2 = e$ (e is called a *projector* of E). The algebra B plays the role of an algebra of functions on a manifold and E is thought of as a module of sections (matter fields) on a vector bundle over this manifold. A (universal) connection or a covariant derivative on E is a map $\nabla : E \to \Omega^1 B \otimes_B E$, where $\Omega^1 B$ denotes the space of (universal) differential 1-forms on B, such that for all $\rho \in \Omega^1 B$, $x \in E$,

$$\nabla(\rho \otimes x) = \rho \nabla(x) + d\rho \otimes x. \tag{1}$$

A different approach to non-commutative gauge theories, based on quantum groups, was introduced in [4]. In this approach one begins with a quantum principal bundle P constructed algebraically as a Hopf-Galois extension of B by a Hopf algebra (quantum group) H (see [13] for a review of Hopf-Galois extensions). One then constructs a vector bundle as a module E associated to B and an H-comodule (corepresentation) V. Any strong connection (gauge field) in P induces then a covariant derivative in E. An example of a quantum principal bundle is the quantum

---

[1] Lloyd's Tercentenary Fellow. On leave from: Department of Theoretical Physics, University of Łódź, Pomorska 149/153, 90–236 Łódź, Poland. E-MAIL: `tb10@york.ac.uk`

[2] E-MAIL: `majid@damtp.cam.ac.uk`

Hopf fibration over the standard quantum two-sphere in [4]. The Dirac q-monopole is the canonical connection on this bundle. In a recent paper [10] the line bundle associated to the q-monopole bundle has been constructed. It has been shown that it is a finitely generated projective module. The covariant derivative induced by the q-monopole connection is the *Grassmannian connection* (see eq. (6) below).

On the other hand, the standard quantum two-sphere is only one of the infinite family of quantum two-spheres constructed in [15]. To describe gauge theory on all such spheres one needs the notions of a *coalgebra principal bundle*, introduced in [5], and an *associated module* as an algebraic version of a coalgebra vector bundle, introduced in [2]. In this note we announce some of the results of a forthcoming paper [6], in which the theory of [5] is generalised and combined with [2] to give the theory of connections on coalgebra principal and associated bundles, illustrated by the example of monopole connection for all quantum two-spheres. The new results of this note then consist of an explicit projective module description of line bundles over all quantum two-spheres, following the method used for the standard quantum sphere in [10].

We work over the field k of real or complex numbers, and we use the standard coalgebra notation. Thus, for a coalgebra C, Δ is the coproduct and ϵ is the counit. In a Hopf algebra S denotes the antipode. We use Sweedler's notation for the coproduct, $\Delta(c) = c_{(1)} \otimes c_{(2)}$ (summation understood). If P is a right C-comodule then $\Delta_P : P \to P \otimes C$ denotes the right coaction. On elements we write $\Delta_P(p) = p_{(0)} \otimes p_{(1)}$ (summation understood). If V is a left C-comodule then ${}_V\Delta : V \to C \otimes V$ denotes the left coaction. The reader not familiar with coalgebras is referred to [16] or to any modern textbook on quantum groups.

COALGEBRA GAUGE THEORY

We begin with the following definition from [3] (which generalises the earlier definition of a coalgebra principal bundle in [5]).

DEFINITION 1 *Let C be a coalgebra, P an algebra and a right C-comodule, and B a subalgebra of P, $B := \{b \in P \mid \forall p \in P\ \Delta_P(bp) = b\Delta_P(p)\}$. We say that P is a* coalgebra-Galois extension *(or a* coalgebra principal bundle*) of B* iff *the canonical left P-module right C-comodule map $can : P \otimes_B P \to P \otimes C$, $can(p \otimes p') = pp'_{(0)} \otimes p'_{(1)}$ is bijective. Such a coalgebra-Galois extension is denoted by $P(B)^C$.*

The definition of a coalgebra principal bundle implies [3] that there exists a unique *entwining map* $\psi : C \otimes P \to P \otimes C$ (cf. [5, Definition 2.1] for a definition of an entwining map). Explicitly, $\psi(c \otimes p) = can \circ (can^{-1}(1 \otimes c)p)$. The properties of ψ [5, Proposition 2.2] imply that the map $\psi^2 = (\mathrm{id}_P \otimes \psi) \circ (\psi \otimes \mathrm{id}_P) : C \otimes P \otimes P \to P \otimes P \otimes C$, restricts to $\psi^2 : C \otimes \Omega^1 P \to \Omega^1 P \otimes C$, where $\Omega^1 P = \{\sum_i p_i \otimes p'_i \in P \otimes P \mid \sum_i p_i p'_i = 0\}$ is the bimodule of universal one-forms on P. Also $\psi^2 \circ (\mathrm{id}_C \otimes d) = (d \otimes \mathrm{id}_C) \circ \psi$, where $d : P \to \Omega^1 P$, $d(p) = 1 \otimes p - p \otimes 1$ is the universal differential. In other words, the universal differential calculus on P is covariant with respect to ψ. This allows one to introduce the following definition in [6], which extends the earlier one in [5].

DEFINITION 2 *Let* $P(B)^C$ *be a coalgebra principal bundle with the entwining map* ψ. *A* connection one-form *in* $P(B)^C$ *is a linear map* $\omega : C \to \Omega^1 P$ *such that:*

(i) $1_{(0)}\omega(1_{(1)}) = 0$, *where* $1_{(0)} \otimes 1_{(1)} = \Delta_P(1)$.

(ii) For all $c \in C$, $\chi \circ \omega(c) = 1 \otimes c - \epsilon(c)\Delta_P(1)$, *where* $\chi : P \otimes P \to P \otimes C$, $\chi(p \otimes p') = p\Delta_P(p')$.

(iii) For all $c \in C$, $\psi^2(c_{(1)} \otimes \omega(c_{(2)})) = \omega(c_{(1)}) \otimes c_{(2)}$.

Conditions (i)-(iii) have the following geometric meaning. By (i), ω can be thought of as acting on $\ker \epsilon$, which can be interpreted as a dual of the "Lie algebra" of C. Condition (ii) means that an application of ω to a vector field obtained by lifting an element of a Lie algebra gives back this element of a Lie algebra. Finally, (iii) is the covariance of ω under the adjoint coaction. As shown in [5] [6], connection 1-forms are in bijective correspondence with *connections*, i.e., projections Π in $\Omega^1 P$ with the fixed kernel $P(\Omega^1 B)P$ and such that $\Pi \circ d$ is right C-covariant. A connection is said to be a *strong connection*, if for all $p \in P$, $dp - p_{(0)}\omega(p_{(1)}) \in (\Omega^1 B)P$ [9].

Given a coalgebra principal bundle $P(B)^C$, and a left C-comodule V with the coaction ${}_V\Delta : V \to C \otimes V$, one constructs the left B-module

$$E := \{\sum_i p^i \otimes v^i \in P \otimes V \mid \sum_i \Delta_P(p_i) \otimes v^i = \sum_i p_i \otimes {}_V\Delta(v^i)\}.$$

The module E, introduced in [2], plays the role of a fibre bundle associated to P with fibre V. The coalgebra gauge theory in [6] ensures that any strong connection ω in P induces a covariant derivative $\nabla : E \to \Omega^1 B \otimes_B E$ on E (in the sense of equation (1)) given by

$$\nabla(\sum_i p^i \otimes v^i) = \sum_i dp^i \otimes v^i - \sum_i {p^i}_{(0)}\omega({p^i}_{(1)}) \otimes v^i. \tag{2}$$

Since the existence of a connection in a module E is equivalent to E being projective [8, Corollary 8.2], we conclude that if there is a strong connection in $P(B)^C$, then every associated left B-module E is projective.

LEFT COALGEBRA LINE BUNDLES ON QUANTUM SPHERES

Recall that the quantum group $SU_q(2)$ is a free algebra generated by 1 and the matrix of generators $\mathbf{t} = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$, subject to the relations:

$$\alpha\beta = q\beta\alpha, \quad \alpha\gamma = q\gamma\alpha, \quad \beta\gamma = \gamma\beta, \quad \beta\delta = q\delta\beta,$$

$$\gamma\delta = q\delta\gamma, \quad \alpha\delta = \delta\alpha + (q - q^{-1})\beta\gamma, \quad \alpha\delta - q\beta\gamma = 1.$$

$SU_q(2)$ is a Hopf algebra of a matrix type, i.e.

$$\Delta(t_{ik}) = \sum_{j=1}^{2} t_{ij} \otimes t_{jk}, \quad \epsilon(\mathbf{t}) = 1, \quad S\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} = \begin{pmatrix} \delta & -q^{-1}\beta \\ -q\gamma & \alpha \end{pmatrix}.$$

The quantum two-spheres S^2_{qs} [15] are homogeneous spaces of $SU_q(2)$. For a given q they can be viewed as subalgebras of $SU_q(2)$ generated by 1 and

$$\xi = s(\alpha^2 - q^{-1}\beta^2) + (s^2-1)q^{-1}\alpha\beta, \quad \eta = s(q\gamma^2 - \delta^2) + (s^2-1)\gamma\delta,$$

$$\zeta = s(q\alpha\gamma - \beta\delta) + (s^2-1)q\beta\gamma,$$

where $s \in [0,1]$. The standard quantum sphere corresponds to $s = 0$[3].

It is shown in [1] that for any quantum sphere there is a coalgebra principal bundle with the base $B = S^2_{qs}$ and the total space $P = SU_q(2)$. The structure coalgebra is a quotient $C_s = SU_q(2)/J_s$ where $J_s = \{\xi - s, \eta + s, \zeta\}SU_q(2)$. We denote by π_s the canonical projection $SU_q(2) \to C_s$. The coproduct on C_s is obtained by projecting down the coproduct in $SU_q(2)$ by π_s. The coaction of C_s on $SU_q(2)$ is given by $\Delta_{SU_q(2)} = (\mathrm{id}_{SU_q(2)} \otimes \pi_s) \circ \Delta$. It has been recently found in [14], that the coalgebras C_s are spanned by group-like elements ($c \in C_s$ is group-like if $\Delta(c) = c \otimes c$). These are computed explicitly in [6] and are given by

$$\pi_s(1), \quad g_n^+ = \pi_s(\prod_{k=0}^{n-1}(\alpha + q^k s\beta)), \quad g_n^- = \pi_s(\prod_{k=0}^{n-1}(\delta - q^{-k}s\gamma)), \quad n = 1, 2, \ldots$$

(all products increase from left to right).

Similarly to [10], consider a one-dimensional left corepresentation $V = k$ of C_s given by the coaction ${}_k\Delta : k \to C_s \otimes k$, ${}_k\Delta(1) = g_1^+ \otimes 1$. The left S^2_{qs}-module E_s associated to $SU_q(2)$ and this corepresentation comes out as

$$E_s = \{x(\alpha + s\beta) + y(\gamma + s\delta) \mid x, y \in S^2_{qs}\}.$$

The module E_s is the *line bundle* over S^2_{qs}. Our first goal is to construct the projector for E_s.

Consider the following matrix with entries from S^2_{qs},

$$e_s = \frac{1}{1+s^2}\begin{pmatrix} 1-\zeta & \xi \\ -\eta & s^2 + q^{-2}\zeta \end{pmatrix}.$$

Notice that e_s can be also written in the following form

$$\begin{aligned} e_s &= \frac{1}{1+s^2}\begin{pmatrix} (\alpha + s\beta)(\delta - qs\gamma) & (\alpha + s\beta)(s\alpha - q^{-1}\beta) \\ (\gamma + s\delta)(\delta - qs\gamma) & (\gamma + s\delta)(s\alpha - q^{-1}\beta) \end{pmatrix} \\ &= \frac{1}{1+s^2}\begin{pmatrix} \alpha + s\beta \\ \gamma + s\delta \end{pmatrix}(\delta - qs\gamma, s\alpha - q^{-1}\beta). \end{aligned}$$

---

[3] It is perhaps more customary to parametrise quantum spheres by the parameter $c \geq 0$ [15]. The parameter s which is used here is related to p in [1] via $s - s^{-1} = p^{-1}$, and p^2 should be identified with c in [15].

The use of the relation

$$(\delta - qs\gamma)(\alpha + s\beta) + (s\alpha - q^{-1}\beta)(\gamma + s\delta) = 1 + s^2, \tag{3}$$

makes it clear that $e_s^2 = e_s$, i.e., e_s is a projector.

Now consider the left S^2_{qs}-module $(S^2_{qs})^2 e_s := \{(x, y)e_s \mid x, y \in S^2_{qs}\}$, and the linear map $\Theta_s : (S^2_{qs})^2 e_s \to E_s$ given by $\Theta_s : (x, y)e_s \mapsto x(\alpha + s\beta) + y(\gamma + s\delta)$. This map is well-defined because $(x, y)e_s = 0$ if and only if

$$\begin{cases} (x(\alpha + s\beta) + y(\gamma + s\delta))(\delta - qs\gamma) = 0 \\ (x(\alpha + s\beta) + y(\gamma + s\delta))(s\alpha - q^{-1}\beta) = 0 \end{cases} \tag{4}$$

Multiplying the first equation by $\alpha + s\beta$ and the second one by $\gamma + s\delta$ and then using (3) one deduces that $x(\alpha + s\beta) + y(\gamma + s\delta) = 0$. The map Θ_s is clearly surjective. Also, if $x(\alpha + s\beta) + y(\gamma + s\delta) = 0$, then (4) are satisfied, so that $(x, y)e_s = 0$. This implies that Θ_s is an isomorphism of left S^2_{qs}-modules. Therefore, E_s is isomorphic to $(S^2_{qs})^2 e_s$ and hence it is a finitely generated projective module.

Next we compute the Chern number or the monopole charge of E_s. The formula (4.4) in [12] for the Chern character of S^2_{qs} gives $\tau^1(1) = 0$, $\tau^1(\zeta) = \frac{1+s^2}{1-q^{-2}}$. Hence the Chern number of E_s is $\mathrm{ch}(E_s) := \tau^1(\mathrm{tr} e_s) = \tau^1(1 - (\zeta - q^{-2}\zeta)/(1 + s^2)) = -1$, and is independent of s. By the same argument as in [10] we conclude that E_s is not isomorphic to $S^2_{qs} \otimes V$. This implies [2] that $SU_q(2)$ is not isomorphic to $S^2_{qs} \otimes C_s$ as a left S^2_{qs}-module and a right C_s-comodule (i.e. $SU_q(2)(S^2_{qs})^{C_s}$ is not *cleft*). This means that the coalgebra Hopf bundle of S^2_{qs} is not trivial.

Finally we show that the *monopole connection* in $SU_q(S^2_{qs})^{C_s}$, constructed in [6], is the Grassmannian connection in E_s. Its connection one-form is given by

$$\omega(g^\pm_n) = Si(g^\pm_n)_{(1)} di(g^\pm_n)_{(2)}, \tag{5}$$

where

$$i(g^+_n) = \prod_{k=0}^{n-1} \frac{\alpha + q^k s(\beta + \gamma) + q^{2k} s^2 \delta}{1 + q^{2k} s^2}, \quad i(g^-_n) = \prod_{k=0}^{n-1} \frac{\delta - q^{-k} s(\beta + \gamma) + q^{-2k} s^2 \alpha}{1 + q^{-2k} s^2}.$$

The connection (5) is strong and the corresponding covariant derivative ∇ on E_s computed from (2) is $\nabla(u) = du - u\omega(g^+_1)$, for all $u \in E_s$. Let $u = x(\alpha + s\beta) + y(\gamma + s\delta)$ for some $x, y \in S^2_{qs}$. Using the explicit form of $\omega(g^+_1)$ we have

$$\begin{aligned} \nabla(u) &= d(x(\alpha + s\beta)) + d(y(\gamma + s\delta)) \\ &\quad - \frac{1}{1 + s^2}(x(\alpha + s\beta) + y(\gamma + s\delta))((\delta - qs\gamma)d(\alpha + s\beta) + (s\alpha - q^{-1}\beta)d(\gamma + s\delta)). \end{aligned}$$

With the help of the Leibniz rule and (3) this can be gathered in the following form

$$\begin{aligned} \nabla(u) &= (d(x, y) + (x, y)\frac{1}{2} d \begin{pmatrix} (\alpha + s\beta)(\delta - qs\gamma) & (\alpha + s\beta)(s\alpha - q^{-1}\beta) \\ (\gamma + s\delta)(\delta - qs\gamma) & (\gamma + s\delta)(s\alpha - q^{-1}\beta) \end{pmatrix}) \begin{pmatrix} \alpha + s\beta \\ \gamma + s\delta \end{pmatrix} \\ &= ((dx, dy) + (x, y)de_s) \begin{pmatrix} \alpha + s\beta \\ \gamma + s\delta \end{pmatrix}. \end{aligned}$$

Now viewing u in $(S^2_{qs})^2 e_s$ via Θ_s^{-1} and ∇ as a map $(S^2_{qs})^2 e_s \to \Omega^1 S^2_{qs} \otimes_{S^2_{qs}} (S^2_{qs})^2 e_s$ via Θ_s one obtains

$$\nabla((x, y)e_s) = (d(x, y) + (x, y)de_s)e_s. \tag{6}$$

The covariant derivative of the form (6) is called the *Grassmannian connection* on a projective module. Therefore, similarly to the standard quantum sphere case discussed in [10], the covariant derivative corresponding to the q-monopole connection on any quantum sphere S^2_{qs} is the Grassmannian connection on E_s.

REFERENCES

1. T. Brzeziński. Quantum homogeneous spaces as quantum quotient spaces. *J. Math. Phys.*, 37:2388–2399, 1996. (q-alg/9509015)
2. T. Brzeziński. On modules associated to coalgebra Galois extensions. *Preprint*, q-alg/9712023
3. T. Brzeziński and P.M. Hajac. Coalgebra extensions and algebra coextensions of Galois type. *Preprint*, q-alg/9708010, 1997 (to appear in *Commun. Algebra*).
4. T. Brzeziński and S. Majid. Quantum group gauge theory on quantum spaces. *Commun. Math. Phys.*, 157:591–638, 1993. Erratum 167:235, 1995.
5. T. Brzeziński and S. Majid. Coalgebra bundles. *Commun. Math. Phys.*, 191:467–492, 1998.
6. T. Brzeziński and S. Majid. Quantum geometry of algebra factorisations and coalgebra bundles. *Preprint* DAMTP/97-139 (in preparation).
7. A. Connes. *Non-Commutative Geometry.* Academic Press, 1994.
8. J. Cuntz and D. Quillen. Algebra extensions and nonsingularity. *J. Amer. Math. Soc.*, 8:251–289, 1995.
9. P. M. Hajac. Strong connections on quantum principal bundles. *Commun. Math. Phys.* 182:579–617, 1996.
10. P.M. Hajac and S. Majid. Projective module description of the q-monopole. *Preprint* math.QA/9803003
11. G. Landi. *An Introduction to Noncommutative Spaces and their Geometries.* Springer-Verlag, 1997.
12. T. Masuda, Y. Nakagami and J. Watanabe. Noncommutative differential geometry on the quantum two sphere of Podleś. I: An algebraic viewpoint. *K-Theory*, 5:151–175, 1991.
13. S. Montgomery. *Hopf Algebras and Their Actions on Rings.* CBMS Lectures vol. 82, AMS, 1993.
14. E.F. Müller and H.-J. Schneider. Quantum homogeneous spaces with faithfully flat module structure. *University of Munich preprint*, 1998.
15. P. Podleś. Quantum spheres. *Lett. Math. Phys.*, 14:193–202, 1987.
16. M. E. Sweedler. *Hopf Algebras* W.A.Benjamin, Inc., New York, 1969.